

k

**Printed by HPS Server
for**

Walk-Up_Printing

Printer: cm1_9e12_gblbptr

Date: 08/10/03

Time: 21:16:02

Document Listing

Document	Selected Pages	Page Range	Copies
WO009305802	47	1 - 47	1
Total (1)	47	-	-



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 37/02, C07K 7/10	A1	(11) International Publication Number: WO 93/05802 (43) International Publication Date: 1 April 1993 (01.04.93)
--	----	--

(21) International Application Number: PCT/US92/07622

(22) International Filing Date: 4 September 1992 (04.09.92)

(30) Priority data:

760,054	13 September 1991 (13.09.91)	US
870,960	20 April 1992 (20.04.92)	US

(71) Applicant: MAGAININ PHARMACEUTICALS INC.
[US/US]; 5110 Campus Drive, Plymouth Meeting, PA 19462 (US).

(72) Inventors: MALOY, W., Lee ; 106 Claremont Drive, Lansdale, PA 19446 (US). KARI, U., Prasad ; 1005 Winfield Court, Lansdale, PA 19446 (US). WILLIAMS, Jon, I. ; 11 Darvel Drive, Robbinsville, NJ 08691 (US).

(74) Agents: OLSTEIN, Elliot, M. et al.; Carella, Byrne, Bain, Gilfillan, Cecchi & Stewart, 6 Becker Farm Road, Roseland, NJ 07068 (US).

(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE).

Published

With international search report.

(54) Title: BIOLOGICALLY ACTIVE AMPHIPHILIC PEPTIDE COMPOSITIONS AND USES THEREFOR

(57) Abstract

Biologically active amphiphilic peptides including one of the following basic structures: I) $R_1-R_2-R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_1-R_1$; II) $R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2$; III) $R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1-R_2-R_3$; IV) $R_1-R_2-R_1-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_4$; V) $R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_3$; or VI) $R_1-R_1-R_3-R_2-R_1-R_1-R_1-R_1-R_1-R_2-R_1-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1$. R_1 is a hydrophobic amino acid and R_2 is a basic hydrophilic or neutral hydrophilic amino acid, R_3 is a neutral hydrophilic amino acid, and R_4 is a neutral hydrophilic amino acid or proline. There is also provided a biologically active amphiphilic peptide having a structural formula selected from the group consisting of: (a) $R_1-R_2-R_2-R_1-R_2-R_2-R_1$; (b) $R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$; (c) $R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$; (d) $R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$; and (e) $R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$, wherein R_1 and R_2 are hereinabove described. Such peptides may be employed as pharmaceuticals.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MN	Mongolia
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Faso	GN	Guinea	NO	Norway
BG	Bulgaria	GR	Greece	NZ	New Zealand
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	PT	Portugal
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovak Republic
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CS	Czechoslovakia	LU	Luxembourg	SU	Soviet Union
CZ	Czech Republic	MC	Monaco	TD	Chad
DE	Germany	MG	Madagascar	TC	Togo
DK	Denmark	ML	Mali	UA	Ukraine
ES	Spain			US	United States of America


-1-

BIOLOGICALLY ACTIVE AMPHIPHILIC PEPTIDE
COMPOSITIONS AND USES THEREFOR

This application is a continuation-in-part of Application Serial No. 760,054, filed September 13, 1991.

This invention relates to biologically active amphiphilic peptides. More particularly, this invention relates to biologically active amphiphilic peptides useful in pharmaceutical compositions.

In accordance with an aspect of the present invention, there is provided a biologically active amphiphilic peptide which includes the following basic structure X_{10} :



R_1 is a hydrophobic amino acid, and R_2 is a basic hydrophilic or neutral hydrophilic amino acid.

In one embodiment, the peptide includes the basic structure $Y_{10}-X_{10}$ wherein X_{10} is as hereinabove described and Y_{10} is:

- (i) R_1 ;
- (ii) R_2-R_1 ; or
- (iii) $R_2-R_2-R_1$, wherein R_1 and R_2 are as hereinabove described.

The hydrophobic amino may be selected from the class consisting of Ala, Cys, Phe, Gly, Ile, Leu, Met, Pro, Val, Trp and Tyr.

SUBSTITUTE SHEET

The basic hydrophilic amino acid acids may be selected from the class consisting of Lys, Arg, His, Orn, homoarginine (Har), 2, 4-diaminobutyric acid (Dbu), and p-aminophenylalanine. The neutral hydrophilic amino acids may be selected from the class consisting of Asn, Gln, Ser, Thr, and homoserine (Hse).

In one embodiment, R_1 is leucine. In another embodiment, R_2 is lysine. Representative examples of peptides in accordance with this aspect of the present invention include those having the following structures:

(SEQ ID NO: 1)

(SEQ ID NO: 2)

(SEQ ID NO: 3)

(SEQ ID NO: 4)

In accordance with another aspect of the present invention, there is provided a biologically active amphiphilic peptide which includes the following basic structure X_{12} :

$R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2$, wherein R_1 is a hydrophobic amino acid and R_2 is a basic hydrophilic or neutral hydrophilic amino acid.

In one embodiment R_1 is leucine. In another embodiment, R_2 is lysine.

In one embodiment, the peptide includes the basic structure $Y_{12}-X_{12}$, wherein X_{12} is as hereinabove described, and Y_{12} is:

- (i) R_2 ;
- (ii) R_1-R_2 ;
- (iii) $R_1-R_1-R_2$;
- (iv) $R_2-R_1-R_1-R_2$; or
- (v) $R_2-R_2-R_1-R_1-R_2$.

In one embodiment, the peptide may have the following structure;

Lys Lys Leu Leu Lys Lys Leu Lys Lys Leu

5

10

Leu Lys Lys Leu Arg Arg

15

(SEQ ID NO.:5)

In another embodiment, the peptide includes the basic structure $X_{12} - Z_{12}$, wherein X_{12} is as hereinabove described, and Z_{12} is:

- (i) R_1 ;
- (ii) $R_1 - R_1$;
- (iii) $R_1 - R_1 - R_2$;
- (iv) $R_1 - R_1 - R_2 - R_2$; or
- (v) $R_1 - R_1 - R_2 - R_2 - R_1$;

In one embodiment, the peptide may have the following structure:

Lys Leu Lys Lys Leu Leu Lys Lys Leu Lys Lys Leu Leu Lys Lys
Leu

5

10

15

(SEQ ID NO:6)

In another embodiment, the peptide may include the structure:

$(Y_{12})_a - X_{12} - (Z_{12})_b$, wherein X_{12} , Y_{12} and Z_{12} are as hereinabove described, and a is 0 or 1, and b is 0 or 1.

In accordance with another aspect of the present invention, there is provided a biologically active amphiphilic peptide which includes the following basic structure X_{14} :

$R_1 - R_2 - R_2 - R_1 - R_1 - R_2 - R_2 - R_1 - R_2 - R_2 - R_1 - R_1 - R_2 - R_2 - R_3$,
wherein R_1 and R_2 are as hereinabove described, and R_3 is a neutral hydrophilic amino acid.

In one embodiment, the peptide may have the following structure:

(SEQ ID NO:7)

In another embodiment, the peptide may have the following structure:

(SEQ ID NO:8)

In accordance with yet another aspect of the present invention, there is provided a biologically active amphiphilic peptide which includes the following basic structure X_{16} :

$R_1-R_2-R_1-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_4$, wherein R_1 and R_2 are as hereinabove described, and R_4 is a neutral hydrophilic amino acid or proline.

In one embodiment, the peptide may include the following structure $Y_{16}-X_{16}$ wherein X_{16} is the basic peptide structure hereinabove described, and Y_{16} is:

- (i) $-R_1$;
- (ii) $-R_1-R_1$;
- (iii) $-R_2-R_1-R_1$;
- (iv) $-R_1-R_2-R_1-R_1$;
- (v) $-R_1-R_1-R_2-R_1-R_1$;
- (vi) $-R_2-R_1-R_1-R_2-R_1-R_1$; or
- (vii) $-R_2-R_2-R_1-R_1-R_2-R_1-R_1$,

wherein R_1 and R_2 are as hereinabove described.

In one embodiment, the peptide may include the structure:

$X_{16}-Z_{16}$, wherein X_{16} is as hereinabove described, and Z_{16} is:

- (i) $-R_2$;
- (ii) $-R_2-R_2$;
- (iii) $-R_2-R_2-R_1$;
- (iv) $-R_2-R_2-R_1-R_1$;
- (v) $-R_2-R_2-R_1-R_1-R_2$;
- (vi) $-R_2-R_2-R_1-R_1-R_2-R_2$; or
- (vii) $-R_2-R_2-R_1-R_1-R_2-R_2-R_1$.

In a preferred embodiment, the peptide may have one of the following structures:

(SEQ ID NO:9); or

(SEQ ID NO:10).

In another embodiment, the peptide may have the structure $(Y_{16})_a-X_{16}-(Z_{16})_b$, wherein X_{16} , Y_{16} , and Z_{16} are as hereinabove described, a is 0 or 1, and b is 0 or 1.

In accordance with another aspect of the present invention, there is provided a biologically active amphiphilic peptide which includes the following basic structure X_{18} :

$R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_3$,

wherein R_1 , R_2 , and R_3 are as hereinabove described.

In accordance with another embodiment, the peptide may include the structure $Y_{18}-X_{18}$, wherein X_{18} is as hereinabove described, and Y_{18} is:

(i) $-R_1$;

(ii) $-R_2-R_1$;

(iii) $-R_2-R_2-R_1$;

(iv) $-R_1-R_2-R_2-R_1$;

(v) $-R_1-R_1-R_2-R_2-R_1$;

(vi) $-R_2-R_1-R_1-R_2-R_2-R_1$; or

(vii) $-R_2-R_2-R_1-R_1-R_2-R_2-R_1$, wherein R_1 and R_2 are as hereinabove described.

In another embodiment, the peptide includes the structure $X_{18}-Z_{18}$, wherein X_{18} is as hereinabove described, and Z_{18} is:

(i) $-R_1$;

(ii) $-R_1-R_5$;

(iii) $-R_1-R_5-R_5$;

(iv) $-R_1-R_5-R_5-R_3$;

(v) $-R_1-R_5-R_5-R_3-R_1$;

(vi) $-R_1-R_5-R_5-R_3-R_1-R_3$;

(vii) $-R_1-R_5-R_5-R_3-R_1-R_3-R_3$;

(viii) $-R_1-R_5-R_5-R_3-R_1-R_3-R_3-R_5$; or

(ix) $-R_1-R_5-R_5-R_3-R_1-R_3-R_3-R_5-R_3$, wherein R_1 and R_3 are as hereinabove described, and R_5 is proline.

In one embodiment, the peptide has the following structure:

(SEQ ID NO:11).

In one embodiment, the peptide may have the structure $(Y_{18})_a-X_{18}-(Z_{18})_b$, wherein X_{18} , Y_{18} , and Z_{18} are as hereinabove described, a is 0 or 1, and b is 0 or 1.

In accordance with another aspect of the present invention, there is provided a biologically active amphiphilic peptide which includes the following basic structure X_{20} :

$R_1-R_1-R_3-R_2-R_1-R_1-R_1-R_1-R_1-R_1-R_2-R_1-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-$
-

wherein R_1 , R_2 , and R_3 are as hereinabove described. In one embodiment, the peptide may have the following structure:

(SEQ ID NO:12).

In another embodiment, the peptide may include the structure $X_{20}-Z_{20}$, wherein X_{20} is as hereinabove described, and Z_{20} is:

- (i) $-R_2$;
- (ii) $-R_2-R_2$;
- (iii) $-R_2-R_2-R_1$;
- (iv) $-R_2-R_2-R_1-R_1$;
- (v) $-R_2-R_2-R_1-R_1-R_2$;
- (vi) $-R_2-R_2-R_1-R_1-R_2-R_2$; or
- (vii) $-R_2-R_2-R_1-R_1-R_2-R_2-R_1$.

In accordance with yet another aspect of the present invention, there is provided a biologically active amphiphilic peptide having a structure selected from the group consisting of:

- (a) $R_1-R_2-R_2-R_1-R_2-R_2-R_1$;
- (b) $R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$;
- (c) $R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$;

- (d) $R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$; and
() $R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$, where in R_1 and R_2 are as hereinabove described.

In one embodiment, the peptide has the structure (a), and a representative example of such a structure is (SEQ ID NO:13), which is given in the accompanying sequence listing.

In another embodiment, the peptide has the structure (b), and a representative example of such a structure is (SEQ ID NO:14), which is given in the accompanying sequence listing.

In another embodiment, the peptide has the structure (c), and a representative example of such a structure is (SEQ ID NO:15) as given in the accompanying sequence listing.

In yet another embodiment, the peptide has the structure (d), and a representative example of such a structure is (SEQ ID NO:16) as given in the accompanying sequence listing.

In a further embodiment, the peptide has the structure (e), and representative examples of such a structure are (SEQ ID NO:17) and (SEQ ID NO:18) as given in the accompanying sequence listing.

In accordance with another aspect of the present invention, there is provided a biologically active amphiphilic peptide having the following structural formula:
(SEQ ID NO:19).

In general, the peptides hereinabove described are ion channel-forming peptides.

An ion channel-forming peptide or ionophore is a peptide which increases the permeability for ions across a natural or synthetic lipid membrane. B. Christensen et al., PNAS, Vol. 85 Pgs. 5072-76 (July, 1988) and Anzai, et al., Biochimica et Biophysica Acta., Vol. 1064, pgs. 256-266 (1991), describe methodology which indicates whether or not

a peptide has ion channel-forming properties and is therefore an ionophore. As used herein an ion channel-forming peptide is a peptide which has ion channel-forming properties as determined by the method of Christensen et al or Anzai, et al.

In one embodiment, the hereinabove described peptides may be acetylated with a CH_3CO -group at the N-terminal, said CH_3CO -group being hereinafter referred to by the letter X.

In another embodiment, the hereinabove described peptides may include an octanoyl group at the N-terminal.

In accordance with one embodiment, each of the amino acid residues contained in the peptides which is not glycine, is a D-amino acid residue. Although the scope of this particular embodiment is not to be limited to any theoretical reasoning, it is believed that the above-mentioned peptides, when consisting entirely of D-amino acid or glycine residues, may have increased resistance to proteolytic enzymes while retaining their biological activity. Such peptides thus may be administered orally. Also, in accordance with another embodiment, all of the amino acid residues may be D-amino acid or glycine residues, or L-amino acid or glycine residues.

In general, the peptides and/or analogues or derivatives thereof are generally water soluble to a concentration of at least 20 mg/ml at neutral pH in water. In addition, the structure of such peptide provides for flexibility of the peptide molecule. When the peptide is placed in water, it does not assume an amphiphilic structure. When the peptide encounters an oily surface or membrane, the peptide chain folds upon itself into a rod-like structure.

The peptides and/or analogues or derivatives thereof may be administered to a host; for example a human or non-human animal, in an amount effective to inhibit growth of a target cell, virus, or virally-infected cell. Thus, for example, the peptides and/or analogues or derivatives thereof may be used as antimicrobial agents anti-viral agents, anti-bacterial agents, anti-tumor agents, anti-parasitic agents, spermicides, as well as exhibiting other bioactive functions.

The term "antimicrobial" as used herein means that the polypeptides of the present invention inhibit, prevent, or destroy the growth or proliferation of microbes such as bacteria, fungi, viruses, or the like.

The term "anti-bacterial" as used herein means that the polypeptides employed in the present invention produce effects adverse to the normal biological functions of bacteria, including death or destruction and prevention of the growth or proliferation of the bacteria when contacted with the polypeptides.

The term "antibiotic" as used herein means that the peptides employed in the present invention produce effects adverse to the normal biological functions of the non-host cell, tissue or organism, including death or destruction and prevention of the growth or proliferation of the non-host cell, tissue, or organism when contacted with the peptides.

The term "spermicidal" as used herein means that the polypeptides employed in the present invention, inhibit, prevent, or destroy the motility of sperm.

The term "antiviral" as used herein means that the polypeptides employed in the present invention inhibit, prevent, or destroy the growth or proliferation of viruses, or of virally-infected cells.

SUBSTITUTE SHEET

The term "anti-tumor" as used herein means that the polypeptide inhibits the growth of or destroys tumors, including cancerous tumors.

The term "anti-parasitic" as used herein means that the polypeptides employed in the present invention inhibit, prevent, or destroy the growth or proliferation of parasites.

The peptides of the present invention have a broad range of potent antibiotic activity against a plurality of microorganisms including gram-positive and gram-negative bacteria, fungi, protozoa, and the like, as well as parasites. The peptides of the present invention allow a method for treating or controlling microbial infection caused by organisms which are sensitive to the peptides. Such treatment may comprise administering to a host organism or tissue susceptible to or afflicted with a microbial infection an antimicrobial amount of at least one of the peptides.

Because of the antibiotic, antimicrobial, antiviral, and antibacterial properties of the peptides, they may also be used as preservatives or sterilants or disinfectants of materials susceptible to microbial or viral contamination.

The peptides and/or derivatives or analogues thereof may be administered in combination with a non-toxic pharmaceutical carrier or vehicle such as a filler, non-toxic buffer, or physiological saline solution. Such pharmaceutical compositions may be used topically or systemically and may be in any suitable form such as a liquid, solid, semi-solid, injectable solution, tablet, ointment, lotion, paste, capsule, or the like. The peptide compositions may also be used in combination with adjuvants, protease inhibitors, or compatible drugs where such a combination is seen to be desirable or advantageous in controlling infection caused by harmful microorganisms including protozoa, viruses, and the like, as well as by parasites.

SUBSTITUTE SHEET

The peptides of the present invention may be administered to a host; in particular a human or non-human animal, in an effective antibiotic and/or anti-tumor and/or anti-viral and/or anti-microbial and/or antibacterial and/or anti-parasitic and/or an antispermicidal amount.

Depending on the use, a composition in accordance with the invention will contain an effective anti-microbial amount and/or an effective antispermicidal amount and/or an effective anti-viral amount and/or an effective anti-tumor amount and/or an effective anti-parasitic and/or an effective antibiotic amount of one or more of the hereinabove described peptides which have such activity. The peptides may be administered by direct application of the peptides to the target cell or virus or virally-infected cell, or indirectly applied through systemic administration.

The peptides of the present invention may also be employed in promoting or stimulating healing of a wound in a host.

The term "wound healing" as used herein includes various aspects of the wound healing process.

These aspects include, but are limited to, increased contraction of the wound, increased deposition of connective tissue, as evidenced by, for example, increased deposition of collagen in the wound, and increased tensile strength of the wound, i.e., the peptides increase wound breaking strength. The peptides of the present invention may also be employed so as to reverse the inhibition of wound healing caused by conditions which depress or compromise the immune system.

The peptides of the present invention may be used in the treatment of external burns and to treat and/or prevent skin and burn infections. In particular, the peptides may be used to treat skin and burn infections caused by

organisms such as, but not limited to, P. aeruginosa and S. aureus.

The peptides are also useful in the prevention or treatment of eye infections. Such infections may be caused by bacteria such as, but not limited to, P. aeruginosa, S. aureus, and N. gonorrhoeae, by fungi such as but not limited to C. albicans and A. fumigatus, by parasites such as but not limited to A. castellani, or by viruses.

The peptides may also be effective in killing cysts, spores, or trophozoites of infection - causing organisms. Such organisms include, but are not limited to Acanthamoeba which forms trophozoites or cysts, C. albicans, which forms spores, and A. fumigatus, which forms spores as well.

The peptides may also be administered to plants in an effective antimicrobial or antiviral or antiparasitic amount to prevent or treat microbial or viral or parasitic contamination thereof.

The peptides, when used in topical compositions, are generally present in an amount of at least 0.1%, by weight. In most cases, it is not necessary to employ the peptide in an amount greater than 2.0%, by weight.

In employing such compositions systemically (intramuscular, intravenous, intraperitoneal), the active peptide is present in an amount to achieve a serum level of the peptide of at least about 5 ug/ml. In general, the serum level of peptide need not exceed 500 ug/ml. A preferred serum level is about 100 ug/ml. Such serum levels may be achieved by incorporating the peptide in a composition to be administered systemically at a dose of from 1 to about 10 mg/kg. In general, the peptide(s) need not be administered at a dose exceeding 100 mg/kg.

The peptides may be produced by known techniques and obtained in substantially pure form. For example, the peptides may be synthesized on an automatic peptide

synthesizer. Journal of the American Chemical Society, Vol. 85, pgs. 2149-54 (1963). It is also possible to produce such peptides by genetic engineering techniques.

Thus, within the scope of the present invention there may be provided DNA which encodes the peptides hereinabove described. Such DNA may be expressed by means known to those skilled in the art. Thus, it is also contemplated that within the scope of the present invention the peptides may be administered by administering to a host DNA which encodes the peptide(s).

In accordance with another embodiment, the peptides of the present invention may be employed in combination with an ion having pharmacological properties for the purposes hereinabove described.

An ion having pharmacological properties is one which when introduced into a target cell, virus, or virally-infected cell, inhibits and/or prevents and/or destroys the growth of the target cell, virus, or virally-infected cell.

Such an ion having pharmacological properties is one which in the absence of an ion channel forming peptide is unable to cross a natural or synthetic lipid membrane; in particular a cell membrane, in sufficient amounts to affect a cell or virus adversely.

The peptide and ion having pharmacological properties may be administered as a single composition or in separate compositions, and the single or separate compositions may include additional materials, actives and/or inactives, in addition to the peptide and ion having pharmacological properties. As representative examples of ions having pharmacological properties which may be employed, there may be mentioned fluoride, peroxide, bicarbonate, silver, zinc, mercury, arsenic, copper, platinum, antimony, gold, thallium, nickel, selenium, bismuth, and cadmium ions.

The peptide and the ion having pharmacological properties, whether administered or prepared in a single composition or in separate compositions, are employed in amounts effective to inhibit and/or prevent and/or destroy the growth of the target cell. In effect, the ion potentiates the action of the peptide, i.e., the amount of ion is effective to reduce the maximum effective concentration of the peptide or protein for inhibiting growth of a target cell, virus, or virally-infected cell.

The ion having pharmacological properties, when used topically, is generally employed in a concentration of from 0.05% to 2.0%. When used systemically, the ion is generally employed in an amount of from 1 to 10 mg. per kg. of host weight. Peptide dosages may be within the ranges hereinabove described.

It is also to be understood that the peptide and ion having pharmacological properties, may be delivered or administered in different forms; for example, the toxic ion may be administered orally, while the peptide may be administered by IV or IP.

As representative examples of administering the peptide or protein and ion for topical or local administration, the peptide could be administered in an amount of up to about 1% weight to weight and the ion delivered in an amount of about 50mM (about 0.1%). Alternatively, the ion, in the form of a salt such as sodium fluoride, could be administered orally in conjunction with systemic administration of the peptide. For example, the peptide may be administered IV or IP to achieve a serum dose of 100 micrograms per milliliter (10 milligrams per kilogram) in conjunction with an oral dose of ion, in particular, sodium fluoride, of 10 meq per kilogram.

In accordance with another embodiment, the peptides of the present invention may be administered to a host in combination with an antibiotic selected from the class

SUBSTITUTE SHEET

consisting of bacitracins, gramacidin, polymyxin, vancomycin, teichoplanin, aminoglycosides, hydrophobic antibiotics, penicillins, monobactams, or derivatives or analogues thereof.

The bacitracins, gramacidin, polymyxin, vancomycin, teichoplanin, and derivatives and analogues thereof, are a group of polypeptide antibiotics. A preferred bacitracin is bacitracin A.

Aminoglycoside antibiotics include tobramycin, kanamycin, amikacin, the gentamicins (e.g., gentamicin C₁, gentamicin C₂, gentamicin C_{1a}), netilmicin, and derivatives and analogues thereof. The preferred aminoglycosides are tobramycin and the gentamicins. The aminoglycosides, and the bacitracins hereinabove described, tend to be hydrophilic and water-soluble.

Penicillins which may be employed include, but are not limited to benzyl penicillin, ampicillin, methicillin (dimethoxyphenyl penicillin), ticarcillin, penicillin V (phenoxymethyl penicillin), oxacillin, cloxacillin, dicloxacillin, flucloxacillin, amoxicillin, and amidinocillin. Preferred penicillins which may be employed are benzyl penicillin and ampicillin. A preferred monobactam which may be employed is aztreonam.

As representative examples of hydrophobic antibiotics which may be used in the present invention, there may be mentioned macrolides such as erythromycin, roxythromycin, clarithromycin, etc.; 9-N-alkyl derivatives of erythromycin; midecamycin acetate; azithromycin; flurithromycin; rifabutin; rokitamycin; a 6-O-methyl erythromycin A known as TE-031 (Taisho); rifapentine; benzypiperazinyl rifamycins such as CGP-7040, CGP-5909, CGP-279353 (Ciba-Geigy); an erythromycin A derivative with a cyclic carbamate fused to the C₁₁/C₁₂ position of a macrolide ring known as A-62514 (Abbott); AC-7230 (Toyo Jozo); benzoxazinorifamycin;

SUBSTITUTE SHEET

difficidin; dirithromycin; a 3-N-piperdinomethylzaino methyl rifamycin SV known as FCE-22250 (Farmitalia); M-119-a (Kirin Brewery); a 6-O-methyl-1-4"-O-carbamoyl erythromycin known as A-63075 (Abbott); 3-formylrifamycin SV-hydrazones with diazabicycloalkyl side chains such as CGP-27557 and CGP-2986 (Ciba-Geigy); and 16-membered macrolides having a 3-O-alpha-L-cladinosyl moiety, such as 3-O-alpha-L-cladinosyldepoxy rosaramicin; tylosins and acyl demycinosyl tylosins.

In addition to the macrolides hereinabove described, rifamycin, carbenicillin, and nafcillin may be employed as well.

Other antibiotics which may be used (whether or not hydrophobic) are antibiotics which are 50-S ribosome inhibitors such as lincomycin; clindamycin; and chloramphenicol; etc.; antibiotics which have a large lipid like lactone ring, such as mystatin; pimaricin, etc.

The peptide and antibiotic may be administered by direct administration to a target cell or by systemic or topical administration to a host which includes the target cell, in order to prevent, destroy or inhibit the growth of a target cell. Target cells whose growth may be prevented, inhibited, or destroyed by the administration of the peptides and antibiotic include Gram-positive and Gram-negative bacteria as well as fungal cells.

The antibiotic, such as those hereinabove described, or derivatives or analogues thereof, when used topically, is generally employed in a concentration of about 0.1% to about 10%. When used systemically, the antibiotic or derivative or analogue thereof is generally employed in an amount of from 1.25 mg. to about 45 mg. per kg. of host weight per day. Peptide dosages may be those as hereinabove described.

As representative examples of administering the peptide and antibiotic for topical or local administration, the

SUBSTITUTE SHEET

peptide could be administered in an amount of from about 0.1% to about 10% weight to weight, and the antibiotic is delivered in an amount of from about 0.1% to about 10% weight to weight.

In accordance with another embodiment, the peptides of the present invention may be administered in combination with an antiparasitic agent or an antifungal agent.

Antiparasitic agents which may be employed include, but are not limited to, anti-protozoan agents. Examples of specific anti-parasitic agents which may be employed include, but are not limited to, pentamidine isethionate, and propamidine isethionate (Brolene).

Anti-fungal agents which may be employed include, but are not limited to, ketoconazole. It is also to be understood that certain anti-parasitic agents, may also have anti-fungal activity, and that certain anti-fungal agents may have anti-parasitic activity.

In accordance with another embodiment, the peptides of the present invention may be administered in combination with an antibiotic which inhibits DNA gyrase, which is an enzyme involved in the formation of bonds between individual coiling strands of replicating bacterial DNA. Thus, DNA gyrase is necessary for the normal replication of bacterial DNA, and, therefore, antibiotics which inhibit DNA gyrase inhibit the normal replication of bacterial DNA.

Examples of antibiotics which inhibit DNA gyrase include nalidixic acid, oxolinic acid, cinoxacin, and quinolone antibiotics which include ciprofloxacin, norfloxacin, ofloxacin, enoxacin, pefloxacin, lomefloxacin, fleroxacin, tosulfloxacin, temafloxacin, and rufloxacin.

In accordance with another embodiment, the peptides of the present invention may be administered for the purpose hereinabove described in combination with other biologically

active amphiphilic peptides, or in combination with ion channel-forming proteins.

The present invention will be further described with respect to the following example; however, the scope of the invention is not to be limited thereby.

EXAMPLE

Table I, which follows, indicates the Minimal Inhibitory Concentration (MIC) in $\mu\text{g/ml}$ of Peptides (SEQ ID NO:1) through (SEQ ID NO:3), (SEQ ID NO:5), (SEQ ID NO:6), (SEQ ID NO:9) through (SEQ ID NO:11), and (SEQ ID NO:17) against S.aureus strain ATCC 25923, P. aeruginosa strain ATCC 27853, and E.coli ATCC strain 25922.

The procedure for the antibacterial assay is based upon the guidelines of the National Committee for Clinical Laboratory Standards, Document M7-T2, Volume 8, No. 8, 1988.

Stock solutions of peptides (SEQ ID NO: 1) through (SEQ ID NO:3), (SEQ ID NO:5), (SEQ ID NO:6), (SEQ ID NO:9) through (SEQ ID NO:12), and (SEQ ID NO:18) in accordance with the present invention are prepared at a concentration of 512 $\mu\text{g/ml}$ in sterile deionized distilled water and stored at -70°C . Each peptide is a C-terminal amide and may or may not be acetylated at the N-Terminus. An acetyl group at the N-terminus is indicated by an X.

The stock peptide solution is diluted in serial dilutions (1:2) down the wells of a microtiter plate so that the final concentrations of peptides in the wells are 0.25, 0.50, 1, 2, 4, 8, 16, 32, 64, 128, and 256 $\mu\text{g/ml}$. $1-5 \times 10^5$ CFUs/ml of either S. aureus ATCC 25923, E. coli ATCC 25922, or P. aeruginosa ATCC 27853 were added to the wells in full strength Mueller Hinton broth (BBL 11443) from a mid-log culture. The inoculum is standardized spectrophotometrically at 600 nm and is verified by colony counts. The plates are incubated for 16-20 hours at 37°C ,

and the minimal inhibitory concentration (MIC) for each peptid is determined. Minimal inhibitory concentration is defined as the lowest concentration of peptide which produces a clear well in the microtiter plate. The minimal inhibitory concentration of each of (SEQ ID NO:1) through (SEQ ID NO:3), (SEQ ID NO:5), (SEQ ID NO:6), (SEQ ID NO:9) through (SEQ ID NO:11), and (SEQ ID NO:17) is given in Table I below.

Table I
Minimal Inhibitory Concentration
($\mu\text{g/ml}$)

<u>Peptide</u>	<u>S.aureus</u>	<u>P.aeruginosa</u>	<u>E.coli</u>
X-(SEQ ID NO:1)-NH ₂	16	4,8	4
(SEQ ID NO.:1)-NH ₂	64,128	8,16	8,16
X-(SEQ ID NO:2)-NH ₂	16	16	8,16
(SEQ ID NO.:2)-NH ₂	16	16	8,16
X-(SEQ ID NO:3)-NH ₂	64	16	8, 16
(SEQ ID NO.:3)-NH ₂	64	16	8,16,32
X-(SEQ ID NO:5)-NH ₂	4	8	32
X-(SEQ ID NO:6)-NH ₂	16	128	8
X-(SEQ ID NO:9)-NH ₂	16	8	64
X-(SEQ ID NO:10)-NH ₂	16,32	16	64
(SEQ ID NO:11)	>256	256	>256
(SEQ ID NO:17)-NH ₂	>256	32,64	64,128
X-(SEQ ID NO:17)-NH ₂	256	8,16	32,64

The peptides of the present invention, whether administered alone or in combination with agents such as ions having pharmacological properties, antibiotics, or other biologically active peptides or proteins as hereinabove described, may be employed in a wide variety of pharmaceutical compositions in combination with a non-toxic pharmaceutical carrier or vehicle such as a filler, non-toxic buffer, or physiological saline solution. Such pharmaceutical compositions may be used topically

or systemically and may be in any suitable form such as a liquid, solid, semi-solid, injectable solution, tablet, ointment, lotion, paste, capsule or the like. The peptide and/or agent as hereinabove described may also be used in combination with adjuvants, protease inhibitors, or compatible drugs where such a combination is seen to be desirable or advantageous in controlling infection caused by harmful microorganisms including protozoa, viruses, parasites, fungi, and the like.

The peptide may be administered to a host in particular an animal, in an effective antibiotic and/or anti-tumor and/or antiviral and/or antimicrobial and/or antispermicidal and/or antifungal and/or antiparasitic amount, or in an amount effective to stimulate wound healing in a host. The peptides may be administered either alone or in combination with an ion having pharmacological properties, antibiotic, or ion channel forming peptide or protein as hereinabove described. When the peptide is administered in combination with a ion having pharmacological properties, the activity of the peptide is potentiated.

When the peptide is administered in combination with an agent as hereinabove described, it is possible to administer the peptide and agent in separate forms. For example, the agent may be administered systemically and the peptide may be administered topically.

When the peptide is administered topically, it may be administered in combination with a water-soluble vehicle, said water-soluble vehicle being in the form of an ointment, cream, lotion, paste or the like. Examples of water-soluble vehicles which may be employed include, but are not limited to, glycols, such as polyethylene glycol, hydroxycellulose, and KY Jelly. The water-soluble vehicle is preferably free of an oily substance.

The peptide may also be employed in combination with a ion having pharmacological properties, as hereinabove described in the form of an oral composition for oral hygiene. Such a composition may be incorporated into a wide variety of compositions

and materials used for oral hygiene purposes, which include, but are not limited to, toothpastes, mouthwashes, tooth gels, and tooth powders. Such composition may thus be used to treat or prevent periodontal disease, to prevent or reduce plaque, and/or to prevent or treat or reduce dental caries. The peptide and toxic ion may be used to inhibit, prevent, or destroy the growth of Streptococcus mutans, which is associated with dental caries and periodontal disease.

Numerous modifications and variations of the present invention are possible in light of the above teachings; therefore, within the scope of the appended claims the invention may be practiced otherwise than as particularly described.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Maloy, W. Lee
Kari, U. Prasad
Williams, Jon I.

(ii) TITLE OF INVENTION: Biologically Active Peptide
Compositions and Uses Therefor

(iii) NUMBER OF SEQUENCES: 19

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Carella, Byrne, Bain, Gilfillan,
Cecchi & Stewart

(B) STREET: 6 Becker Farm Road

(C) CITY: Roseland

(D) STATE: New Jersey

(E) COUNTRY: USA

(F) ZIP: 07068

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: 3.5 inch diskette

(B) COMPUTER: IBM PS/2

(C) OPERATING SYSTEM: PC-DOS

(D) SOFTWARE: DW4.V2

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

-23-

(A) APPLICATION NUMBER: 07/760,054

(B) FILING DATE: 13-SEP-1991

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Olstein, Elliot M.

(B) REGISTRATION NUMBER: 24,025

(C) REFERENCE/DOCKET NUMBER: 421250-161

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 201-994-1700

(B) TELEFAX: 201-994-1744

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 11 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Leu Lys Lys Leu Lys Lys Leu Leu Lys Leu

5

10

Leu

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 12 amino acids

SUBSTITUTE SHEET

- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Leu Leu Lys Lys Leu Lys Lys Leu Leu Lys
 5 10
Leu Leu

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:3:

Lys Leu Leu Lys Lys Leu Lys Lys Leu L u

5

10

Lys Leu Leu

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:4:

Lys Lys Leu Leu Lys Lys Leu Lys Lys Leu

5

10

Leu Lys Leu Leu

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

SUBSTITUTE SHEET

(D) OTHER INFORMATION: May be a C-terminal amide, and/or may be acetylated at N-terminus.

(ix) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Lys Lys Leu Leu Lys Lys Leu Lys Lys Leu Leu Lys Lys Leu Arg Arg
5 10 15

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Lys Leu Lys Lys Leu Leu Lys Lys Leu Lys
5 10
Lys Leu Leu Lys Leu Leu
15

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:7:

Leu Lys Lys Leu Leu Lys Lys Leu Lys Lys
 5 10
Leu Leu Lys Lys Asn
 15

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus, Xaa is homoserine.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:8:

Leu Lys Lys Leu Leu Lys Lys Leu Lys Lys
 5 10
Leu Leu Lys Lys Xaa
 15

SUBSTITUTE SHEET

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Leu Lys Leu Leu Lys Lys Leu Leu Lys Lys
5 10
Asn Lys Lys Leu Leu Lys Lys Leu
15

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

SUBSTITUTE SHEET

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:10:

Leu Lys Leu L u Lys Lys Leu Leu Lys Lys
 5 10
 Pro Lys Lys Leu Leu Lys Lys Leu
 15

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:11:

Leu Leu Lys Lys Leu Lys Lys Leu Leu Lys
 5 10
 Lys Leu Gln Gly Pro Pro Gln Gly Gln Ser
 15 20
 Pro Gln

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:

SUBSTITUTE SHEET

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:12:

Leu	Ala	Ser	Lys	Ala	Gly	Ala	Ile	Ala	Gly
			5					10	
Lys	Ile	Ala	Lys	Lys	Leu	Leu	Lys	Lys	Leu
			15					20	

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:13:

Leu	Lys	Lys	Leu	Lys	Lys	Leu
			5			

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:14:

Leu Leu Lys Lys Leu Lys Lys Leu

5

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

UBSTITUTE SHEET

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:15:

Lys Leu Leu Lys Lys Leu Lys Lys Leu

5

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:16:

Lys Lys Leu Leu Lys Lys Leu Lys Lys Leu

5

10

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(D) OTHER INFORMATION: May be a C-terminal amide,

SUBSTITUTE SHEET

and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:17:
Leu Lys Lys Leu Leu Lys Lys Leu Lys Lys Leu
5 10

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:18:
Ala Lys Lys Leu Leu Lys Lys Leu Lys Lys Leu
5 10

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(D) OTHER INFORMATION: May be a C-terminal amide,
and/or may be acetylated at N-terminus.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Leu Lys Lys Leu Leu Lys Lys Leu Lys Lys Leu

5

10

Leu Lys Arg

WHAT IS CLAIMED IS

1. A biologically active amphiphilic peptide, said peptide including the following basic structure X_{10} :

$R_1-R_2-R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_1-R_1$, wherein R_1 is a hydrophobic amino acid, and R_2 is a basic hydrophilic or neutral hydrophilic amino acid.

2. The peptide of Claim 1 wherein the peptide includes the basic structure:

$Y_{10}-X_{10}$, wherein X_{10} is the basic peptide structure of Claim 1, and Y_{10} is:

- (i) R_1 ;
- (ii) R_2-R_1 ; or
- (iii) $R_2-R_2-R_1$.

3. A biologically active amphiphilic peptide, said peptide including the following basic structure X_{12} :

$R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2$, wherein R_1 is a hydrophobic amino acid and R_2 is a basic hydrophilic or neutral hydrophilic amino acid.

4. The peptide of Claim 3 wherein said peptide includes the basic structure $Y_{12}-X_{12}$, wherein X_{12} is the basic peptide structure of Claim 3, and Y_{12} is:

- (i) R_2 ;
- (ii) R_1-R_2 ;
- (iii) $R_1-R_1-R_2$;
- (iv) $R_2-R_1-R_1-R_2$; or
- (v) $R_2-R_2-R_1-R_1-R_2$.

5. The peptide of Claim 3 wherein said peptide includes the basic structure $X_{12}-Z_{12}$, wherein X_{12} is the basic peptide structure of Claim 3, and Z_{12} is:

- (i) R_1 ;
- (ii) R_1-R_1 ;
- (iii) $R_1-R_1-R_2$;
- (iv) $R_1-R_1-R_2-R_2$; or
- (v) $R_1-R_1-R_2-R_2-R_1$.

6. A process for inhibiting growth of a target cell or virus in a host, comprising:

administering to a host a biologically active amphiphilic peptide, said peptide including the following basic structure X_{10} :

$R_1-R_2-R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_1-R_1$, wherein R_1 is a hydrophobic amino acid, and R_2 is a basic hydrophilic or neutral hydrophilic amino acid, said peptide being administered in an amount effective to inhibit growth of a target cell or virus in a host.

7. The process of Claim 6 wherein said peptide includes the basic structure:

$Y_{10}-X_{10}$, wherein X_{10} is the basic peptide structure of Claim 4 and Y_{10} is:

- (i) R_1 ;
- (ii) R_2-R_1 ; or
- (iii) $R_2-R_2-R_1$.

8. The process of Claim 6 wherein said administering is in an effective antimicrobial amount.

9. The process of Claim 6 wherein said administering is in an effective anti-viral amount.

10. The process of Claim 6 wherein said administering is in an effective anti-tumor amount.

11. The process of Claim 6 wherein said administering is in an effective anti-parasitic amount.

12. The process of Claim 6 wherein said administering is in an effective antibiotic amount.

13. A process for inhibiting growth of a target cell or virus in a host, comprising:

administering to a host a biologically active amphiphilic peptide, said peptide including the following basic peptide structure X_{12} :

$R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2$, wherein R_1 is a hydrophobic amino acid and R_2 is a basic hydrophilic or neutral

hydrophilic amino acid, said peptide being administered in an amount effective to inhibit growth of a target cell or virus in a host.

14. The process of Claim 13 wherein the peptide includes the basic structure $Y_{12}-X_{12}$, wherein X_{12} is the basic peptide structure of Claim 13 and Y_{12} is:

- (i) R_2 ;
- (ii) R_1-R_2 ;
- (iii) $R_1-R_1-R_2$;
- (iv) $R_2-R_1-R_1-R_2$; or
- (v) $R_2-R_2-R_1-R_1-R_2$.

15. The process of Claim 13 wherein the peptide includes the basic structure $X_{12}-Z_{12}$, wherein X_{12} is the basic structure of Claim 13, and Z_{12} is:

- (i) R_1 -;
- (ii) R_1-R_1 ;
- (iii) $R_1-R_1-R_2$;
- (iv) $R_1-R_1-R_2-R_2$; or
- (v) $R_1-R_1-R_2-R_2-R_1$.

16. The process of Claim 13 wherein said peptide is administered in an effective antimicrobial amount.


17. The process of Claim 13 wherein said peptide is administered in an effective anti-viral amount.

18. The process of Claim 13 wherein said peptide is administered in an effective anti-tumor amount.

19. The process of Claim 13 wherein said peptide is administered in an effective antibiotic amount.

20. The process of Claim 13 wherein said peptide is administered in an effective anti-parasitic amount.

21. A biologically active amphiphilic peptide, said peptide including the following basic structure X_{14} :



wherein R_1 is a hydrophobic amino acid, R_2 is a basic hydrophilic or neutral hydrophilic amino acid, and R_3 is a neutral hydrophilic amino acid.

22. A biologically active amphiphilic peptide, said peptide including the following basic structure X_{16} :

$R_1-R_2-R_1-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_4$, wherein R_1 is a hydrophobic amino acid, R_2 is a basic hydrophilic or a neutral hydrophilic amino acid, and R_4 is a neutral hydrophilic amino acid or proline.

23. The peptide of Claim 22 wherein said peptide includes the basic structure $X_{16}-Z_{16}$, wherein X_{16} is the basic peptide structure of Claim 22, and Z_{16} is:

- (i) $-R_2$;
- (ii) $-R_2-R_2$;
- (iii) $-R_2-R_2-R_1$;
- (iv) $-R_2-R_2-R_1-R_1$;
- (v) $-R_2-R_2-R_1-R_1-R_2$;
- (vi) $-R_2-R_2-R_1-R_1-R_2-R_2$; or
- (vii) $R_2-R_2-R_1-R_1-R_2-R_2-R_1$.

24. A biologically active amphiphilic peptide, said peptide including the following basic peptide structure X_{18} :

$R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_3$, wherein R_1 is a hydrophobic amino acid, R_2 is a basic hydrophilic or neutral hydrophilic amino acid, and R_3 is a neutral hydrophilic amino acid.

25. The peptide of Claim 24 wherein the peptide includes the basic structure $X_{18}-Z_{18}$, wherein X_{18} is the basic peptide structure of Claim 24, and Z_{18} is:

- (i) $-R_1$;
- (ii) $-R_1-R_5$;
- (iii) $-R_1-R_5-R_5$;
- (iv) $-R_1-R_5-R_5-R_3$;
- (v) $-R_1-R_5-R_5-R_3-R_1$;
- (vi) $-R_1-R_5-R_5-R_3-R_1-R_3$;

- (vii) $-R_1-R_5-R_5-R_3-R_1-R_3-R_3$;
 (viii) $-R_1-R_5-R_5-R_3-R_1-R_3-R_3-R_5$; or
 (ix) $-R_1-R_5-R_5-R_3-R_1-R_3-R_3-R_5-R_3$, wherein R_1 is a hydrophobic amino acid, R_3 is a neutral hydrophilic amino acid, and R_5 is proline.

26. A biologically active amphiphilic peptide, said peptide including the following basic structure X_{20} :

$R_1-R_1-R_3-R_2-R_1-R_1-R_1-R_1-R_1-R_2-R_1-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-$,
 wherein R_1 is a hydrophobic amino acid, R_2 is a basic hydrophilic or neutral hydrophilic amino acid, and R_3 is a neutral hydrophilic amino acid.

27. A biologically active amphiphilic peptide having a structure selected from the group consisting of:

- (a) $R_1-R_2-R_2-R_1-R_2-R_2-R_1$;
 (b) $R_1-R_1-R_2-R_2-R_1-R_2-R_1-R_1$;
 (c) $R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$;
 (d) $R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$; and
 (e) $R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1$, wherein R_1 is a hydrophobic amino acid, and R_2 is a basic hydrophilic or neutral hydrophilic amino acid.

28. A biologically active amphiphilic peptide having the following structural formula:

(SEQ ID NO:19).

29. A process for inhibiting growth of a target cell, virus, or virally-infected cell in a host, comprising:

administering to a host a biologically active amphiphilic peptide, said peptide including the following basic structure X_{14} :

$R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_3$,
 wherein R_1 is a hydrophobic amino acid, R_2 is a basic hydrophilic or neutral hydrophilic amino acid, and R_3 is a neutral hydrophilic amino acid, said peptide being administered in an amount effective to inhibit growth of a target cell, virus, or virally-infected cell in a host.

30. The process of Claim 29 wherein said administering is in an effective antimicrobial amount.

31. The process of Claim 29 wherein said administering is in an effective anti-viral amount.

32. The process of Claim 29 wherein said administering is in an effective anti-tumor amount.

33. The process of Claim 29 wherein said administering is in an effective anti-parasitic amount.

34. The process of Claim 29 wherein said administering is in an effective antibiotic amount.

35. A process for inhibiting growth of a target cell, virus, or virally-infected cell in a host, comprising:

administering to a host a biologically active amphiphilic peptide, said peptide including the following basic structure X_{16} :

$R_1-R_2-R_1-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_4$, wherein R_1 is a hydrophobic amino acid, R_2 is a basic hydrophilic or a neutral hydrophilic amino acid, and R_4 is a neutral hydrophilic amino acid or proline, said peptide being administered in an amount effective to inhibit growth of a target cell, virus, or virally-infected cell in a host.

36. The process of Claim 35 wherein said peptide includes the basic structure $X_{16}-Z_{16}$, wherein X_{16} is the basic peptide structure of Claim 35, and Z_{16} is:

- (i) $-R_2$;
- (ii) $-R_2-R_2$;
- (iii) $-R_2-R_2-R_1$;
- (iv) $-R_2-R_2-R_1-R_1$;
- (v) $-R_2-R_2-R_1-R_1-R_2$;
- (vi) $-R_2-R_2-R_1-R_1-R_2-R_2$; or
- (vii) $R_2-R_2-R_1-R_1-R_2-R_2-R_1$.

37. The process of Claim 35 wherein said administering is in an effective antimicrobial amount.

SUBSTITUTE SHEET

38. The process of Claim 35 wherein said administering is in an effective anti-viral amount.

39. The process of Claim 35 wherein said administering is in an effective anti-tumor amount.

40. The process of Claim 35 wherein said administering is in an effective anti-parasitic amount.

41. The process of Claim 35 wherein said administering is in an effective antibiotic amount.

42. A process for inhibiting growth of a target cell, virus, or virally-infected cell in a host, comprising:

administering to a host a biologically active amphiphilic peptide, said peptide including the following basic structure X_{18} :

$R_1-R_1-R_2-R_2-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_3$,

wherein R_1 is a hydrophobic amino acid, R_2 is a basic hydrophilic or neutral hydrophilic amino acid, and R_3 is a neutral hydrophilic amino acid, said peptide being administered in an amount effective to inhibit growth of a target cell, virus, or virally-infected cell in a host.

43. The process of Claim 42 wherein the peptide includes the basic structure $X_{18}-Z_{18}$, wherein X_{18} is the basic peptide structure of Claim 42, and Z_{18} is:

- (i) $-R_1$;
- (ii) $-R_1-R_5$;
- (iii) $-R_1-R_5-R_5$;
- (iv) $-R_1-R_5-R_5-R_3$;
- (v) $-R_1-R_5-R_5-R_3-R_1$;
- (vi) $-R_1-R_5-R_5-R_3-R_1-R_3$;
- (vii) $-R_1-R_5-R_5-R_3-R_1-R_3-R_3$;
- (viii) $-R_1-R_5-R_5-R_3-R_1-R_3-R_3-R_5$; or
- (ix) $-R_1-R_5-R_5-R_3-R_1-R_3-R_3-R_5-R_3$, wherein R_1 is a hydrophobic amino acid, R_3 is a neutral hydrophilic amino acid, and R_5 is proline.

SUBSTITUTE SHEET

44. The process of Claim 42 wherein said administering is in an effective antimicrobial amount.

45. The process of Claim 42 wherein said administering is in an effective anti-viral amount.

46. The process of Claim 42 wherein said administering is in an effective anti-tumor amount.

47. The process of Claim 42 wherein said administering is in an effective anti-parasitic amount.

48. The process of Claim 42 wherein said administering is in an effective antibiotic amount.

49. A process for inhibiting growth of a target cell, virus, or virally-infected cell in a host, comprising:

administering to a host a biologically active amphiphilic peptide, said peptide including the following basic structure X_{20} :

$R_1-R_1-R_3-R_2-R_1-R_1-R_1-R_1-R_1-R_2-R_1-R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-$,
wherein R_1 is a hydrophobic amino acid, R_2 is a basic hydrophilic or neutral hydrophilic amino acid, and R_3 is a neutral hydrophilic amino acid, said peptide being administered in an amount effective to inhibit growth of a target cell, virus, or virally-infected cell in a host.

50. The process of Claim 49 wherein said administering is in an effective antimicrobial amount.

51. The process of Claim 49 wherein said administering is in an effective anti-viral amount.

52. The process of Claim 49 wherein said administering is in an effective anti-tumor amount.

53. The process of Claim 49 wherein said administering is in an effective anti-parasitic amount.

54. The process of Claim 49 wherein said administering is in an effective antibiotic amount.

55. A process for inhibiting growth of a target cell, virus, or virally-infected cell in a host comprising:

SUBSTITUTE SHEET

administering to a host a biologically active amphiphilic peptide having a structure selected from the group consisting of:

- (a) $R_1-R_2-R_2-R_1-R_2-R_2-R_1$;
- (b) $R_1-R_1-R_2-R_2-R_1-R_2-R_1$;
- (c) $R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_1$;
- (d) $R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_1$; and
- (e) $R_1-R_2-R_2-R_1-R_1-R_2-R_2-R_1-R_2-R_1$, wherein R_1 is a hydrophobic amino acid, and R_2 is a basic hydrophilic or neutral hydrophilic amino acid, said peptide being administered in an amount effective to inhibit growth of a target cell, virus, or virally-infected cell in a host.

56. The process of Claim 55 wherein said administering is in an effective antimicrobial amount.

57. The process of Claim 55 wherein said administering is in an effective anti-viral amount.

58. The process of Claim 55 wherein said administering is in an effective anti-tumor amount.

59. The process of Claim 55 wherein said administering is in an effective anti-parasitic amount.

60. The process of Claim 55 wherein said administering is in an effective antibiotic amount.

61. A process for inhibiting growth of a target cell, virus, or virally-infected cell in a host, comprising:

administering to a host a biologically active amphiphilic peptide having the following structural formula:

(SEQ ID NO:19), said peptide being administered in an amount effective to inhibit growth of a target cell, virus, or virally-infected cell in a host.

62. The process of Claim 61 wherein said administering is in an effective antimicrobial amount.

63. The process of Claim 61 wherein said administering is in an effective anti-viral amount.

64. The process of Claim 61 wherein said administering is in an effective anti-tumor amount.

65. The process of Claim 61 wherein said administering is in an effective anti-parasitic amount.

66. The process of Claim 61 wherein said administering is in an effective antibiotic amount.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US92/07622**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(5) :A61K 37/02; C07K 7/10

US CL :514/12 530/324

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/12 530/324

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO, A, 89/04371 (Jaynes et al) 18 May 1989, see entire document.	1-66
Y	Peptide Chemistry, issued 1986, Mihara et al., "Design and Synthesis of Basic Amphiphilic Peptides Possessing Antimicrobial Activity and Their Interaction with Lipids, pages 223-228, see entire document.	1-66

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be part of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 DECEMBER 1992

Date of mailing of the international search report:

28 DEC 1992

Name and mailing address of the ISA/
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

LESTER L. LEE

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-0196

HPS Trailer Page
for
Walk-Up_Printing

UserID: k

Printer: cm1_9e12_gblbptr

Summary

<u>Document</u>	<u>Pages</u>	<u>Printed</u>	<u>Missed</u>	<u>Copies</u>
WO009305802	47	47	0	1
Total (1)	47	47	0	-